



**Human Health Risk
Assessment:**
Host Technology Energy
from Waste, Hownsgill
Industrial Park

October 2020



Experts in air quality
management & assessment

Document Control

Client	Project Genesis	Principal Contact	Lee Searles (Enzygo Ltd)
---------------	-----------------	--------------------------	--------------------------

Job Number	J4203
-------------------	-------

Report Prepared By:	Dr Imogen Heard
----------------------------	-----------------

Document Status and Review Schedule

Report No.	Date	Status	Reviewed by
J4203A/3/F1	23 October 2020	Final	Dr Ben Marner (Technical Director)

This report has been prepared by Air Quality Consultants Ltd on behalf of the Client, taking into account the agreed scope of works. Unless otherwise agreed, this document and all other Intellectual Property Rights remain the property of Air Quality Consultants Ltd.

In preparing this report, Air Quality Consultants Ltd has exercised all reasonable skill and care, taking into account the objectives and the agreed scope of works. Air Quality Consultants Ltd does not accept any liability in negligence for any matters arising outside of the agreed scope of works. The Company operates a formal Quality Management System, which is certified to ISO 9001:2008, and a formal Environmental Management System, certified to ISO 14001:2004. QMF 08.

When issued in electronic format, Air Quality Consultants Ltd does not accept any responsibility for any unauthorised changes made by others.

When printed by Air Quality Consultants Ltd, this report will be on Evolve Office, 100% Recycled paper.

Air Quality Consultants Ltd
23 Coldharbour Road, Bristol BS6 7JT Tel: 0117 974 1086
12 Airedale Road, London SW12 8SF Tel: 0208 673 4313
aqc@aqconsultants.co.uk

Registered Office: 12 St Oswalds Road, Bristol, BS6 7HT
 Companies House Registration No: 2814570

Executive Summary

- 1.1 A human health risk assessment has been carried out in order to identify any potential health risks associated with emissions of dioxins and furans from the waste to power facility at Hownsgill Technology Park in Consett, County Durham. The assessment draws on the dispersion modelling presented in the Air Quality Assessment prepared to support the planning application, and additionally uses the United States Environmental Protection Agency's Human Health Risk Assessment Protocol (HHRAP). A range of worst-case assumptions has been applied and thus the assessment is very precautionary.
- 1.2 The assessment has shown that the risks to health comply with the relevant benchmarks. All of the impacts are assessed to be insignificant. No specific mitigation measures are required beyond those incorporated into the design of the facility.

Contents

1	Introduction	3
2	Scope.....	4
3	Assessment Approach	5
4	Assessment Criteria	12
5	Results	15
6	Conclusions	22
7	References.....	23
8	Glossary.....	24
9	Appendices	26
A1	Professional Experience.....	27

Tables

Table 1:	Assumed Dioxin and Furan Emissions from the proposed facility	6
Table 2:	Receptors Used in the Health Risk Assessment.....	8
Table 3:	Site-specific Parameters Used	10
Table 4:	Reference Dose and Hazard Quotient.....	12
Table 5:	Tolerable Daily Intake (TDI) Values	13
Table 6:	Soil Guideline Values (SGVs) for Dioxins and Furans ^a	14
Table 7:	Total Lifetime Cancer Risk for all Receptors ^a	15
Table 8:	Total Lifetime Carcinogenic Risk by Pathway at Worst-case Receptors ^a	16
Table 9:	Hazard Quotient for All Receptors ^a	16
Table 10:	Hazard Quotient by Pathway at Worst-case Receptors ^a	17
Table 11:	Predicted Oral Intake of Dioxins and Furans at Worst-case Receptors (pg-TEQ/kg/d).....	19
Table 12:	Predicted Oral Intake of Dioxins and Furans at Worst-case Receptors (pg-TEQ/kg/week)	20
Table 13:	Estimated Infant Average Daily Dose (ADD) of Dioxins, furans and PCBs (pg-TEQ'kg/d).....	21
Table 14:	Maximum Process Contribution to Dioxin and Furan Concentrations in Soils..	21

Figures

Figure 1:	Receptors for Health Risk Assessment	9
-----------	--	---

1 Introduction

- 1.1 A detailed human health risk assessment has been carried out in order to identify any potential health risks associated with emissions from the waste to power facility (“the proposed facility”) at Hownsgill Technology Park in Consett, County Durham. The assessment has been carried out by Air Quality Consultants Ltd on behalf of Project Genesis.
- 1.2 The pollutants of concern to this assessment (as defined in paragraph 2.1) are termed ‘persistent’, which means that they have the potential to remain and accumulate in the local environment over the operational life of the facility, including bioaccumulation through the food chain. The assessment is based on the United States Environmental Protection Agency (USEPA) Human Health Risk Assessment Protocol (HHRAP). This approach is current best-practice in the UK and is often recommended by the UK Environment Agency. The assessment has used the Industrial Risk Assessment Program-Human Health (IRAP) model, which is based on the HHRAP.
- 1.3 The HHRAP method was developed to assess risks in areas where much of the population derives a substantial portion of its diet from local produce. Because of the potential for accumulation in the food chain, animal husbandry (particularly the raising of pigs and cattle) forms a key part of HHRAP. In practice, in many areas near to the proposed facility the HHRAP method will provide an extremely conservative (i.e. worst-case) assessment.

2 Scope

- 2.1 Following standard Environment Agency practice for studies of this nature, the health risk assessment focuses on the uptake of polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans, often abbreviated to 'dioxins and furans'. While other substances such as metals, acid gases, and particulate matter would be emitted from the proposed facility, these can be adequately assessed by comparing the predicted ambient concentrations against relevant assessment criteria. A detailed assessment of these other pollutants is presented in the Air Quality Assessment prepared to support the planning application.
- 2.2 Exposure to emissions of dioxins and furans can be through a number of pathways, with inhalation and the food chain being most critical. This assessment covers exposure through the direct inhalation of dioxins and furans as gases and fine airborne particles, as well as indirect exposure following the deposition of contaminants to land and subsequent transfer by biogeochemical processes through soils and vegetation into the food chain.
- 2.3 The assessment has not considered the potential for significant impacts associated with ingestion of drinking water, or dermal contact with soil or water. Other studies have shown that absorption through dermal contact with soil is significantly less efficient than absorption by inhalation and can thus safely be discounted (Pasternach, 1989). Dermal contact with standing water in the area will also be sporadic and unlikely to lead to significant exposure. Exposure via drinking water requires contamination of drinking water sources local to the point of consumption. Mains water for the local area comes from reservoirs throughout the Northumbrian Water region. The nearest reservoir is located approximately 7 km north west of the proposed facility. As the prevailing winds in the area are predominantly from the west, the reservoir will only be downwind of the development site infrequently. At a distance of 7 km, it is not expected that mains water will be significantly affected by emissions from the proposed facility and this pathway has thus not been considered further in this assessment.
- 2.4 The assessment has also discounted the potential for local residents to derive a large proportion of their diet from locally-caught fish. Any fish caught in the local area are unlikely to provide a substantial part of people's diets, therefore this pathway has not been included in this assessment.

3 Assessment Approach

Baseline Conditions

- 3.1 Background exposure to dioxins and furans is dependent on a wide range of complex individual factors and will vary from one individual to another. Key issues include individual lifestyle, diet, baseline land quality, and background levels of dioxins and furans. This study has thus focused on the potential impact of the proposed facility, rather than on the combined effect of the proposed facility and other potential sources of exposure. If it is possible to demonstrate that the impacts of the proposed facility will be insignificant, then it is not necessary to calculate the total exposure.

Emissions of Dioxins and Furans from the Proposed Facility

- 3.2 The general terms 'dioxins and furans' describe a large number of compounds, which are individually known as 'congeners'. Each congener has a different toxicity and physical properties with regard to atmospheric behaviour.
- 3.3 As explained in the Air Quality Assessment prepared to support the planning application, the Industrial Emissions Directive (IED) (2010/75/EU, 2010) allows a maximum emission rate for total dioxins and furans of 0.1 ng/Nm³ (averaged over a sampling period of between 6 and 8 hours). The IED also provides a range of congener-specific international toxic equivalence (TEQ) factors by which the mass concentrations of specific congeners should be multiplied before summing. In other words, 0.1 ng/Nm³ refers not to the total mass emission, but to the toxic-equivalent emission. In practice, the proposed facility is unlikely to emit anywhere near 0.1 ng/Nm³, but it has nevertheless been assumed that 0.1 ng TEQ/Nm³ would be emitted during every hour of every year. This provides a worst-case assessment.
- 3.4 Her Majesty's Inspectorate of Pollution (HMIP) previously published a dioxin and furan congener profile based on measurements at municipal waste incinerators (HMIP, 1996). This provides the most robust dataset to describe the proposed facility at this time and has thus been used to define the relative quantities of each congener that would be emitted.
- 3.5 The congener profile from HMIP has been combined with the IED toxic equivalence factor and the volumetric emission rate from the proposed facility to calculate g/s mass emissions of the individual congeners. These calculations are shown in Table 1.

Table 1: Assumed Dioxin and Furan Emissions from the proposed facility

Congener	Sum of Emissions (ng-TEQ/Nm ³)	% of Total Toxic-Equivalent Conc from Each Congener	Toxic Eq Factor (ng TEQ / ng)	Mass Emission Conc (ng/Nm ³) ^a	Mass Emission Rate (g/s) ^b
2,3,7,8-TCDD	-	3.1%	1	0.0031	2.94 x 10 ⁻¹¹
1,2,3,7,8-PeCDD	-	12.4%	0.5	0.025	2.37 x 10 ⁻¹⁰
1,2,3,4,7,8-HxCDD	-	2.9%	0.1	0.029	2.75 x 10 ⁻¹⁰
1,2,3,7,8,9-HxCDD	-	2.1%	0.1	0.021	1.99 x 10 ⁻¹⁰
1,2,3,6,7,8-HxCDD	-	2.6%	0.1	0.026	2.47 x 10 ⁻¹⁰
1,2,3,4,6,7,8-HpCDD	-	1.7%	0.01	0.17	1.61 x 10 ⁻⁹
OCDD	-	0.4%	0.001	0.4	3.80 x 10 ⁻⁹
2,3,7,8-TCDF	-	2.7%	0.1	0.027	2.56 x 10 ⁻¹⁰
2,3,4,7,8-PeCDF	-	26.8%	0.5	0.054	5.13 x 10 ⁻¹⁰
1,2,3,7,8-PeCDF	-	1.4%	0.05	0.028	2.66 x 10 ⁻¹⁰
1,2,3,4,7,8-HxCDF	-	21.8%	0.1	0.22	2.09 x 10 ⁻⁹
1,2,3,7,8,9-HxCDF	-	0.4%	0.1	0.0042	3.99 x 10 ⁻¹¹
1,2,3,6,7,8-HxCDF	-	8.0%	0.1	0.081	7.69 x 10 ⁻¹⁰
2,3,4,6,7,8-HxCDF	-	8.6%	0.1	0.087	8.26 x 10 ⁻¹⁰
1,2,3,4,6,7,8-HpCDF	-	4.4%	0.01	0.44	4.18 x 10 ⁻⁹
1,2,3,4,7,8,9-HpCDF	-	0.4%	0.01	0.043	4.08 x 10 ⁻¹⁰
OCDF	-	0.4%	0.001	0.36	3.42 x 10 ⁻⁹
Sum	0.1	100%	-	2.1	1.92 x 10⁻⁸

^a i.e. 0.1 ng TEQ/Nm³ divided by the toxic equivalence factor and multiplied by the percentage of total toxic-equivalent concentration from each congener. Calculated from unrounded numbers.

^b i.e. mass emission concentration (ng/Nm³) divided by 1 billion, multiplied by the volume flow rate (34,175 Nm³/hr – see the Air Quality Assessment) divided by 3,600.

Dispersion Modelling

- 3.6 The transport of emissions through air has been modelled using the ADMS-5 dispersion model. The model, and the input parameters used for this assessment, are described in the Air Quality Assessment prepared to support the planning application.
- 3.7 Dry and wet deposition of particle-bound and vapour-phase congeners has been simulated within the ADMS-5 model. In terms of the dry deposition of particles, because large particles will be filtered out from the stack emissions, particulate emissions are expected to be predominantly in the 1-2 µm

diameter range. For particles of this size, deposition velocities are likely to be in the range of 0.001 m/s to 0.1 m/s. For dry deposition of particle-bound congeners, a fixed worst-case deposition velocity of 0.01 m/s has thus been assumed. The dry deposition of vapour-phase congeners has been modelled using the ADMS default parameters for non-reactive gases.

- 3.8 The ADMS model has been run using five separate yearly meteorological datasets (2015 to 2019 inclusive) from the Albemarle meteorological monitoring site. These are the data used in the Air Quality Assessment and are described therein. For each receptor, the maximum predicted concentration, dry deposition flux, and wet deposition flux during any year was taken and used for the health risk assessment. Where these maxima come from different years, the approach is worst-case since, in practice, one form of deposition depletes the potential for the other.

Receptors for Dispersion Modelling

- 3.9 Modelling has been carried out across a 10 km x 10 km model domain, with the stack at the centre. Concentrations have been predicted over this area using nested Cartesian grids. These grids have a spacing of 5 m x 5 m within 200 m of the facility, 25 m x 25 m within 1 km of the facility, 50 m x 50 m within 2 km of the facility, 250 m x 250 m within 2.5 km of the facility, and 500 m x 500 m within 5 km of the facility. The gridded receptors have been modelled at a height of 1.5 m, to represent ground-level human exposure.

Human Health Risk Assessment

- 3.10 The assessment has used the IRAP model, which is based on HHRAP.

Receptors for IRAP Model

- 3.11 Within the IRAP model, there are four receptor types of relevance to this assessment:

- resident adult;
- resident child;
- farmer adult; and
- farmer child.

- 3.12 Resident receptors are assumed to have the potential to intake pollutants via inhalation, by eating above-ground home-grown vegetables and by eating soil¹.

- 3.13 Farmer receptors are assumed to intake pollutants by these same pathways, but also from eating home-reared beef, chicken and pork, drinking milk from cows kept at home, and eating home-produced eggs. It is important to recognise that, when a receptor is included as a farmer, the assumption is made that the location is an active farm at which only beef, pork, poultry, eggs, milk

¹ This is usually accidental and associated with home-grown vegetables.

and vegetables produced at that farm are consumed. It is unlikely that there are any such locations in the vicinity of the proposed facility.

- 3.14 Occasionally, residents keep animals in domestic gardens. Where this occurs, it is highly unlikely that these animals would provide the dominant source of consumed food. While it is possible that, for example, chickens kept at home could provide the only source of eggs consumed, it is relatively unlikely that a dominant proportion of the chickens' food would be grown within the study area.
- 3.15 The results from the ADMS-5 dispersion model runs for the Cartesian grid of receptors have been plotted on a map and used to select the worst-case receptors for the human health risk assessment (i.e. the human health risk assessment receptors were chosen to represent locations where the gridded concentrations and deposition fluxes were highest). The receptors used for the health risk assessment are described in Table 2 and shown in Figure 1.

Table 2: Receptors Used in the Health Risk Assessment

Receptor	Description	Type
A	Farmland north west of the development	Farm
B	Farmland east of the development	Farm
C	Residential property east of the development	Residential
D	Residential property north east of the development	Residential
E	Farmland north of the development	Farm
F	Farmland west of the development	Farm
G	Farmland south of the development	Farm
H	Farmland south west of the development	Farm
I	Residential property north west of the development	Residential
J	Residential property north west of the development	Residential
K	Residential property north of the development	Residential
L	Residential property south east of the development	Residential
M	Farmland west of the development	Farm
N	Residential property west of the development	Residential




Figure 1: Receptors for Health Risk Assessment

Contains Ordnance Survey data © Crown copyright and database right 2020. Ordnance Survey licence number 100046099. Additional data sourced from third parties, including public sector information licensed under the Open Government Licence v1.0.

IRAP Model Inputs

- 3.16 The outputs from the ADMS-5 dispersion model have been used as inputs to the IRAP model. From these, IRAP calculated the concentrations of the various congeners in the environmental media, foodstuffs, and the human population.
- 3.17 The IRAP model requires a wide range of input data to be defined as set out below. In many cases the default parameters have been used, as is common practice for such assessments.

Physical and Chemical Properties of the Pollutants

- 3.18 These parameters determine how the pollutants behave in the environment and their presence and accumulation in various food sources. The HHRAP and IRAP default values have been used in this assessment.

Site-specific Information

- 3.19 The IRAP model requires information about the location and its surroundings including:

- the fraction of animal feed grown on local soils and the amount of animal feed and soil consumed by various species;
- the interception fraction for above ground vegetation, forage and silage and lengths of vegetation exposure to deposition;
- data for assessing the risks with exposure to breast milk, such as infant bodyweight, fraction of mother's body weight that is fat etc.; and
- other physical parameters such as soil and air density etc.

The HHRAP and IRAP default parameters have been used for all of this information.

3.20 Other parameters need to be taken from site-specific data. These are set out in Table 3.

Table 3: Site-specific Parameters Used

Parameter	Value
Annual Mean Precipitation	75.2 cm/yr ^a
Annual Mean Wind Velocity	3.2 m/s ^b
Annual Mean Evapotranspiration	52.7 cm/yr ^c
Annual Mean Irrigation	2.64 cm/yr
Annual Mean Runoff	7.5 cm/yr ^d
Time period over which deposition occurs	30 yrs

^a The average of five year's meteorological data.

^b The average of five year's meteorological data.

^c Assumed to be 70% of precipitation.

^d Assumed to be 10% of precipitation.

Receptor Information

3.21 For each receptor type, IRAP requires information on receptor behaviour including absolute and relative food consumption rates, body weight, inhalation rates, and exposure frequency etc. The HHRAP and IRAP default values have been used in this assessment.

IRAP Model Outputs

Cancer Risk

3.22 The HHRAP approach is to calculate the probability of affected receptors developing cancer over their lifetime as a result of emissions from the proposed facility. This risk is presented as an absolute number which represents a probability. For example, a value of 1×10^{-6} (or 0.000001) represents a one in one million chance of an effect over a lifetime's exposure.

Hazard Risk

- 3.23 IRAP has also been used to calculate the 'Hazard Quotient' for each receptor. The 'Hazard Quotient' is a way of expressing the ratio of the predicted exposure level and a 'reference dose' which represents the level at which no adverse effects are expected. Hazard Quotients are pollutant and pathway specific and can then be summed to calculate a 'Hazard Index'. Of the congeners considered in this assessment, a reference dose has only been set for the most toxic congener (2,3,7,8-TCDD) via the ingestion pathway. Thus, for this assessment, the Hazard Quotient and the Hazard Index are the same. The Hazard Quotient is simply the exposure level divided by the reference dose and is thus unitless.

Oral Intake of All Congeners

- 3.24 While the Hazard Quotient only takes account of intake of the most toxic congener (2,3,7,8-TCDD), the total oral intake of all modelled congeners has been derived from the IRAP model and compared against defined 'Tolerable Daily Intake Levels' (TDIs). The units for this assessment are toxic-equivalent-picogrammes² per kg of bodyweight per day (pg-TEQ/kg/d).

Infant Exposure through Breast Milk

- 3.25 Dioxins and furans are extremely fat soluble and hence prone to accumulation in breast milk. The accumulated compounds can then be passed to the infant. Exposure through breast milk is measured by the Average Daily Dose (ADD) on the basis of an averaging time of one year. The units for this assessment are also pg-TEQ/kg/d.

Concentrations in Soils

- 3.26 The final set of outputs derived from the IRAP model is the concentrations of dioxins and furans in soils. These are expressed as the sum of all congeners in µg per kg (µg/kg) of soil (dry weight).

² One picogramme is equal to 0.000000000001 grammes.

4 Assessment Criteria

Cancer Risk

- 4.1 One definition of ‘acceptable risk’ that has been widely used in the UK, is if exposure to a substance increases a person’s chance of dying in any one year by one chance in a million (1:1,000,000 or 1×10^{-6}) or less³ (Hunter and Fewtrell, 2010). The one in one million figure used in the UK is an annual figure. Since risk is cumulative, assuming a 70 year lifetime, the equivalent lifetime risk is one in fourteen thousand (1:14,300 or 7×10^{-5}).
- 4.2 HHRAP uses a value of one in one hundred thousand (1:100,000 or 1×10^{-5}) for lifetime cancer risk. This is effectively more stringent than the 1 in 1 million annual risk figure and has thus been used for this study. The assessment criterion for cancer risk is thus 1×10^{-5} .

Hazard Risk

- 4.3 As explained in Paragraph 3.23, the Hazard Quotient is the predicted exposure level divided by the ‘reference dose’. The HHRAP reference dose relevant to this study is given in Table 4. If the Hazard Quotient is less than 1 (i.e. the predicted exposure is less than the reference dose), then no adverse health effects are expected. If the Hazard Quotient is greater than 1, then adverse health effects are possible. The Hazard Quotient is different from the cancer risk estimates; in that it cannot be translated to a probability that adverse health effects will occur and it is unlikely to be proportional to risk. It is important to note that a Hazard Quotient exceeding 1 does not necessarily mean that adverse effects will occur; it simply indicates the potential for an effect.

Table 4: Reference Dose and Hazard Quotient

Description	Value
Reference Dose (of 2,3,7,8-TCDD per kg body weight) (mg/kg/d)	1×10^{-9}
Hazard Quotient	1

Oral Intake of All Congeners

- 4.4 The World Health Organisation (WHO) has recommended a range of TDIs (Table 5). These represent a lifetime’s exposure; short-term exceedances of the TDI are not of concern if the long-term average is not exceeded (WHO, 1998).
- 4.5 In the UK, the Health Protection Agency and the Department of Health are advised by the Committee on the Toxicity of Chemicals in Food, Consumer Products and the Environment (COT). The COT has also provided recommendations on TDIs, which are based on adverse effects on a developing

³ By way of comparison, the likelihood of dying in a road traffic accident is approximately 1 in 17,500 per year <http://www.medicine.ox.ac.uk/bandolier/booth/Risk/transportpop.html>

foetus (since this represents the most sensitive exposure pathway) (COT, 2001). The TDIs used in this study are set out in Table 5.

Table 5: Tolerable Daily Intake (TDI) Values

Reference Source	TDI (pg-TEQ/kg/d)
(WHO, 1998)	1 – 4
(COT, 2001)	2

- 4.6 The European Food Safety Authority's (EFSA) expert Panel on Contaminants in the Food Chain (CONTAM) has more recently set a tolerable weekly intake (TWI) of 2 pg-TEQ/kg/week (EFSA, 2018), which has also been accepted by the COT and is also used in this study.

Infant Exposure through Breast Milk

- 4.7 There is no official UK or USEPA assessment criterion for acceptable infant exposure. One approach that is often taken is to compare the ADD against average background exposure levels, while other studies have cited a threshold value of 50 pg-TEQ/kg/d of the congener 2,3,7,8-TCDD as being potentially harmful.
- 4.8 Two separate approaches have been taken here. The first is to compare the ADD for all congeners against an assumed nominal baseline dose of 100 pg-TEQ/kg/day⁴. The second is to compare the ADD for the congener 2,3,7,8-TCDD against the threshold value of 50 pg-TEQ/kg/day.

Concentrations in Soils

- 4.9 The Environment Agency has developed Soil Guideline Values (SGVs) for dioxins and furans (Environment Agency, 2009). These are assessment criteria that can be used to evaluate the risk to human health from long-term exposure to chemicals in soil.
- 4.10 For residential and allotment land uses, the SGVs are based on estimates representative of exposure of young children because they are generally more likely to have higher exposures to soil contaminants. The toxic equivalence of the various congeners has been taken into account in the derivation of the SGVs. Thus, the mass sum of all congeners in the soil may be compared directly with the SGVs.

⁴ A compilation of European dioxin exposure data (DETR, 1999) found that, in 1993, the dioxin intake of firstborn infants, up to 2 months of age, might be around 106 pg-TEQ/kg/day in rural areas of the EU and 144 pg TEQ/kg/day in industrial areas. These intake levels fall rapidly after the first few months. These are well in excess of the TDIs, but, as explained in Paragraph 3.5, the TDIs refer to an average lifetime's exposure. Exposure is likely to have fallen since 1993 but these values nevertheless provide a basis for assessment.

- 4.11 The SGVs used in this assessment are set out in Table 6. These are based on a generic profile of congeners which is representative of general diffuse pollution, but within the context of this assessment are considered to be sufficiently accurate.

Table 6: Soil Guideline Values (SGVs) for Dioxins and Furans^a

Land use	Sum of Congeners ($\mu\text{g}/\text{kg DW}$)
Residential	8
Allotment	8
Commercial	240

^a Based on an assumed soil profile for urban soils and on a sandy loam soil.

Generic Environment Agency Screening Criteria

- 4.12 The Environment Agency has considered potential impacts from industrial emissions in its guidance (Environment Agency, 2016). This explains that regardless of what the baseline environmental conditions are, a process can be considered as insignificant if the long-term (annual mean) process contribution is <1% of the long-term environmental standard.
- 4.13 The Environment Agency has considered potential impacts from industrial emissions in its guidance (Environment Agency, 2016). This explains that regardless of what the baseline environmental conditions are, a process can be considered as insignificant if the long-term (annual mean) process contribution is <1% of the long-term environmental standard.
- 4.14 It should be recognised that this criterion determines when an impact can be screened out as insignificant. It does not imply that impacts will necessarily be significant above this level merely that above this level there is a potential for significant impacts.

5 Results

Cancer Risk

- 5.1 The total lifetime cancer risks associated with emissions from the facility for each of the receptors are presented in Table 7.
- 5.2 The data in Table 7 show that the highest residential risk is predicted at receptor C, where the predicted lifetime risk of developing cancer as a result of emissions from the proposed facility is 6.8×10^{-9} (1 in 147 million), while the highest risk at any of the receptors that have been treated as farms is at receptor B, where the risk is 2.7×10^{-6} (1 in 370 thousand). Receptors B and C are both to the east of the proposed facility. All of the results are well below the assessment criterion of 1×10^{-5} (1 in one hundred thousand). Such risks are conventionally considered to be acceptable in the UK.

Table 7: Total Lifetime Cancer Risk for all Receptors ^a

R	Description	Type ^b	Child	Adult
A	Farmland north west of the development	Fa	1.5×10^{-8}	7.1×10^{-8}
B	Farmland east of the development	Fa	5.8×10^{-7}	2.7×10^{-6}
C	Residential property east of the development	R	3.6×10^{-9}	6.8×10^{-9}
D	Residential property north east of the development	R	1.3×10^{-9}	2.5×10^{-9}
E	Farmland north of the development	Fa	3.8×10^{-8}	1.5×10^{-7}
F	Farmland west of the development	Fa	3.1×10^{-8}	2.8×10^{-7}
G	Farmland south of the development	Fa	2.5×10^{-8}	1.2×10^{-7}
H	Farmland south west of the development	Fa	7.0×10^{-8}	3.3×10^{-7}
I	Residential property north west of the development	R	4.4×10^{-10}	8.0×10^{-10}
J	Residential property north west of the development	R	4.8×10^{-10}	8.6×10^{-10}
K	Residential property north of the development	R	2.2×10^{-10}	4.0×10^{-10}
L	Residential property south east of the development	R	1.6×10^{-10}	2.8×10^{-10}
M	Farmland west of the development	Fa	4.2×10^{-8}	2.0×10^{-7}
N	Residential property west of the development	R	3.4×10^{-10}	6.2×10^{-10}
Assessment Criterion			1×10^{-5}	

1×10^{-5} = 1 in one hundred thousand chance of an effect over a lifetime's exposure

1×10^{-6} = 1 in one million chance of an effect over a lifetime's exposure

1×10^{-8} = 1 in one hundred million chance of an effect over a lifetime's exposure

1×10^{-10} = 1 in one billion chance of an effect over a lifetime's exposure

^a The highest relevant values are shown in bold.

^b R = residential, Fa = farmer.

5.3 Table 8 shows the pathways of exposure for receptors B and C. The most significant pathway for the residential receptors is eating home-grown vegetables and the most significant pathway for farm receptors is drinking home-produced milk. As explained in Paragraph 3.13, by including receptors as farms, it is assumed that the location is an active farm at which only beef, pork, poultry, eggs, milk and vegetables produced at that farm are consumed. As there are unlikely to be any such locations in the vicinity of the proposed facility (i.e. all local residents who consume these foodstuffs are likely to purchase at least some of their food), the assessment is worst-case.

Table 8: Total Lifetime Carcinogenic Risk by Pathway at Worst-case Receptors ^a

Pathway	Receptor B (Farmer)		Receptor C (Resident)	
	Child	Adult	Child	Adult
Drinking home-produced milk	5.2 x 10⁻⁷	2.1 x 10⁻⁶	0	0
Eating home-reared beef	5.4 x 10 ⁻⁸	5.9 x 10 ⁻⁷	0	0
Eating above ground vegetables	5.1 x 10 ⁻⁹	1.5 x 10 ⁻⁸	2.9 x 10⁻⁹	6.2 x 10⁻⁹
Eating home-reared pork	2.8 x 10 ⁻⁹	2.5 x 10 ⁻⁸	0	0
Eating soil	7.8 x 10 ⁻¹⁰	6.3 x 10 ⁻¹⁰	6.2 x 10 ⁻¹⁰	3.3 x 10 ⁻¹⁰
Eating home-reared chicken	1.1 x 10 ⁻¹¹	1.2 x 10 ⁻¹⁰	0	0
Eating home-produced eggs	7.8 x 10 ⁻¹²	8.1 x 10 ⁻¹¹	0	0
Inhalation	5.8 x 10 ⁻¹¹	3.8 x 10 ⁻¹⁰	4.7 x 10 ⁻¹¹	2.3 x 10 ⁻¹⁰
Total Lifetime Risk	5.8 x 10⁻⁷	2.7 x 10⁻⁶	3.6 x 10⁻⁹	6.8 x 10⁻⁹

1 x 10⁻⁵ = 1 in one hundred thousand chance of an effect over a lifetime's exposure

1 x 10⁻⁶ = 1 in one million chance of an effect over a lifetime's exposure

1 x 10⁻⁸ = 1 in one hundred million chance of an effect over a lifetime's exposure

1 x 10⁻¹⁰ = 1 in one billion chance of an effect over a lifetime's exposure

^a Highest relevant values in each column are shown in bold.

Hazard Risk

5.4 The Hazard Quotient for each of the receptors is set out in Table 9. All of the values are well below 1 and the risk of significant health effects is thus discounted. The Hazard Risk pathway breakdown for receptors B and C is given in Table 10. Again, this shows that eating home-grown vegetables is the dominant risk pathway for residents, while for farmers it is drinking home-produced milk.

Table 9: Hazard Quotient for All Receptors ^a

R	Description	Type ^b	Child	Adult
A	Farmland north west of the development	Fa	4.3 x 10 ⁻⁵	2.9 x 10 ⁻⁵
B	Farmland east of the development	Fa	2.8 x 10⁻³	1.9 x 10⁻³

R	Description	Type ^b	Child	Adult
C	Residential property east of the development	R	1.1 x 10⁻⁵	4.4 x 10⁻⁶
D	Residential property north east of the development	R	4.2 x 10 ⁻⁶	1.6 x 10 ⁻⁶
E	Farmland north of the development	Fa	1.8 x 10 ⁻⁴	1.2 x 10 ⁻⁴
F	Farmland west of the development	Fa	2.7 x 10 ⁻⁴	1.9 x 10 ⁻⁴
G	Farmland south of the development	Fa	1.1 x 10 ⁻⁴	7.8 x 10 ⁻⁵
H	Farmland south west of the development	Fa	3.3 x 10 ⁻⁴	2.3 x 10 ⁻⁴
I	Residential property north west of the development	R	1.4 x 10 ⁻⁶	5.0 x 10 ⁻⁷
J	Residential property north west of the development	R	1.5 x 10 ⁻⁶	5.3 x 10 ⁻⁷
K	Residential property north of the development	R	6.8 x 10 ⁻⁷	2.6 x 10 ⁻⁷
L	Residential property south east of the development	R	4.8 x 10 ⁻⁷	1.8 x 10 ⁻⁷
M	Farmland west of the development	Fa	2.0 x 10 ⁻⁴	1.4 x 10 ⁻⁴
N	Residential property west of the development	R	1.1 x 10 ⁻⁶	3.9 x 10 ⁻⁷
Assessment Criterion			1	

$$1 \times 10^{-3} = 0.001$$

$$1 \times 10^{-6} = 0.000001$$

^a Highest relevant values are shown in bold.

^b R = residential, F = farm.

Table 10: Hazard Quotient by Pathway at Worst-case Receptors^a

Pathway	Receptor B (Farmer)		Receptor C (Resident)	
	Child	Adult	Child	Adult
Drinking home-produced milk	2.5 x 10⁻³	1.5 x 10⁻³	0	0
Eating home-reared beef	2.6 x 10 ⁻⁴	4.2 x 10 ⁻⁴	0	0
Eating home-reared pork	1.3 x 10 ⁻⁵	1.7 x 10 ⁻⁵	0	0
Eating above ground vegetables	1.8 x 10 ⁻⁵	7.4 x 10 ⁻⁶	1.0 x 10⁻⁵	4.2 x 10⁻⁶
Eating soil	1.7 x 10 ⁻⁶	1.8 x 10 ⁻⁷	1.3 x 10 ⁻⁶	1.4 x 10 ⁻⁷
Eating home-reared chicken	2.5 x 10 ⁻⁸	3.6 x 10 ⁻⁸	0	0
Inhalation	3.3 x 10 ⁻⁸	3.3 x 10 ⁻⁸	2.7 x 10 ⁻⁸	2.7 x 10 ⁻⁸
Eating home-produced eggs	1.7 x 10 ⁻⁸	2.4 x 10 ⁻⁸	0	0

$$1 \times 10^{-3} = 0.001$$

$$1 \times 10^{-6} = 0.000001$$

$$1 \times 10^{-9} = 0.000000001$$

^a Highest relevant values in each column shown in bold.

Oral Intake of all Congeners

- 5.5 Table 11 sets out the predicted oral intake of all congeners for receptors B and C. The results are separately compared with the TDIs from both WHO and COT. The predicted intakes at receptor C are well below 1% of the TDIs. The predicted intakes at receptor B exceed 1% of both TDIs, predominantly as a result of drinking home-produced milk. It should be recognised that exceedance of the assessment criterion does not necessarily mean that impacts will be significant, merely that there is a potential for significant impacts and that further consideration is required, as set out below.
- 5.6 The impacts at receptor B, as shown in Table 11, are up to 4.0% of the lower-bound WHO assessment criterion and up to 2.0% of the COT assessment criterion. While it is not possible to immediately discount these changes as insignificant, they remain small. It must also be recognised that this assessment is very much worst-case. The figures in Table 11 are based on the assumption that the proposed facility will emit continuously at the maximum emission rates allowed under the IED. Furthermore, as discussed in paragraph 3.13, when a receptor is included as a farmer, the assumption is made that the location is an active farm at which only beef, pork, poultry, eggs, milk and vegetables produced at that farm are consumed. It is highly unlikely that receptor E meets these conditions, as it represents the worst-case point within an individual field close to the proposed facility. Taking these factors into account, it is considered that the predicted oral intake presented in Table 11 represents an extreme worst case. The actual impacts are expected to be much lower and are therefore judged to be insignificant.

Table 11: Predicted Oral Intake of Dioxins and Furans at Worst-case Receptors (pg-TEQ/kg/d)

Pathway	Receptor B (Farmer)		Receptor C (Resident)	
	Child	Adult	Child	Adult
Drinking home-produced milk	3.5×10^{-2}	2.1×10^{-2}	0	0
Eating home-reared beef	3.7×10^{-3}	6.0×10^{-3}	0	0
Eating home-reared pork	2.0×10^{-4}	2.7×10^{-4}	0	0
Eating above ground vegetables	4.1×10^{-4}	1.7×10^{-4}	2.4×10^{-4}	1.0×10^{-4}
Eating soil	1.0×10^{-4}	1.1×10^{-5}	8.0×10^{-5}	8.6×10^{-6}
Eating home-reared chicken	1.4×10^{-6}	2.1×10^{-6}	0	0
Eating home-produced eggs	9.7×10^{-7}	1.4×10^{-6}	0	0
Total Intake	4.0×10^{-2}	2.8×10^{-2}	3.2×10^{-4}	1.1×10^{-4}
WHO Assessment Criteria (TDI)	1-4			
COT Assessment Criteria (TDI)	2			
Intake as % of 1 pg-TEQ/kg/d	4.0	2.8	0.03	0.01
Intake as % of 2 pg-TEQ/kg/d	2.0	1.4	0.02	0.006

$$1 \times 10^{-3} = 0.001$$

$$1 \times 10^{-6} = 0.000001$$

$$1 \times 10^{-9} = 0.000000001$$

5.7 Table 12 sets out the predicted oral intake of all congeners for receptors B and C and compares the results with the TWI from the EFSA. The predicted intake at receptor C is well below 1% of the TWI. The predicted intake at receptor B exceeds 1% of the TWI, again predominantly as a result of drinking home-produced milk.

5.8 As stated in paragraph 5.6, the assessment is very much worst-case. The figures in Table 12 are based on the assumption that the proposed proposed facility will emit continuously at the maximum emission rates allowed under the IED. When a receptor is included as a farmer, the assumption is made that the location is an active farm at which only beef, pork, poultry, eggs, milk and vegetables produced at that farm are consumed. Taking these factors into account, it is considered that the predicted oral intake presented in Table 12 represents an extreme worst case. The actual impacts are expected to be much lower and are therefore judged to be insignificant.

Table 12: Predicted Oral Intake of Dioxins and Furans at Worst-case Receptors (pg-TEQ/kg/week)

Pathway	Receptor B (Farmer)		Receptor C (Resident)	
	Child	Adult	Child	Adult
Drinking home-produced milk	2.5×10^{-1}	1.5×10^{-1}	0	0
Eating home-reared beef	2.6×10^{-2}	4.2×10^{-2}	0	0
Eating home-reared pork	1.4×10^{-3}	1.9×10^{-3}	0	0
Eating above ground vegetables	2.9×10^{-3}	1.2×10^{-3}	1.7×10^{-3}	7.1×10^{-4}
Eating soil	7.1×10^{-4}	7.6×10^{-5}	5.6×10^{-4}	6.0×10^{-5}
Eating home-reared chicken	9.9×10^{-6}	1.5×10^{-5}	0	0
Eating home-produced eggs	6.8×10^{-6}	9.5×10^{-6}	0	0
Total Intake	2.8×10^{-1}	1.9×10^{-1}	2.2×10^{-3}	7.7×10^{-4}
EFSA Assessment Criteria (TWI)	2			
Intake as % of 2 pg-TEQ/kg/week	13.9	9.7	0.1	0.04

$$1 \times 10^{-3} = 0.001$$

$$1 \times 10^{-6} = 0.000001$$

Infant Exposure through Breast Milk

- 5.9 Table 13 sets out the estimated ADDs for infant exposure through breast milk for receptors B and C. The predicted ADDs have been compared with the assessment criteria described in Paragraph 4.8. All of the ADDs are less than 1% of the respective criteria and the impacts can thus be discounted as insignificant.

Table 13: Estimated Infant Average Daily Dose (ADD) of Dioxins, furans and PCBs (pg-TEQ'kg/d)

Pathway	Receptor B (Farmer)	Receptor C (Resident)
All Congeners		
ADD (pg TEQ/kg/day)	0.41	0.002
Assessment Criterion (pg TEQ/kg/day)	100	
ADD as percentage of Criterion	0.41	0.002
2,3,7,8-TCDD		
ADD (pg TEQ/kg/day)	0.06	0.0002
Assessment Criterion (pg TEQ/kg/day)	50	
ADD as percentage of Criterion	0.12	0.0005

Concentrations in Soils

- 5.10 Table 14 sets out the maximum process contributions to dioxin and furan concentrations in soils at receptors B and C. The predicted concentration has been compared with the residential SGV defined in Table 6. The predicted concentrations are well below 1% of the SGV. The impacts can thus be discounted as insignificant.

Table 14: Maximum Process Contribution to Dioxin and Furan Concentrations in Soils

Parameter	Receptor B (Farmer)	Receptor C (Resident)
Process Contribution to Maximum Soil Concentration ($\mu\text{g}/\text{kg}$)	0.0002	0.0001
Soil Guideline Value (SGV) ($\mu\text{g}/\text{kg}$)	8	
Process Contribution as % of Guideline Value	0.002	0.002

6 Conclusions

6.1 The human health risk assessment has shown that:

- **cancer risk** - all of the predicted process contributions are less than the assessment criterion. They are at levels which are conventionally considered to be acceptable and are thus discounted as **insignificant**;
- **hazard risk** - all of the predicted process contributions are less than 1 and the risk of effects can be discounted as **insignificant**;
- **oral intake of all congeners** - all of the predicted process contributions at relevant locations are small and judged to be **insignificant**;
- **infant exposure through breast milk** -all of the predicted process contributions are less than 1% of the relevant assessment criteria and can thus be discounted as **insignificant**; and
- **concentrations in soils** - all of the predicted process contributions are less than 1% of the relevant assessment criterion and can thus be discounted as **insignificant**.

6.2 These conclusions are on the basis that the proposed facility will emit dioxins and furans constantly at the maximum level permitted by the European Directive (2010/75/EU, 2010). In practice, emissions are expected to be a small fraction of those assessed; meaning that the impacts will have been grossly over-stated.

6.3 This assessment has found no requirement for specific mitigation measures beyond those incorporated into the design of the facility.

7 References

2010/75/EU, D. (2010). *Directive on Industrial Emissions*.

COT. (2001). *COT statement on the tolerable daily intake for dioxins and dioxin-like polychlorinated biphenyls*. Retrieved from <http://cot.food.gov.uk/committee/committee-on-toxicity/cotstatements/cotstatementsyrs/cotstatements2001/dioxinsstate>

Defra. (2009). *Land Contamination: using soil guidance values (SGVs)*.

DETR. (1999). *Compilation of EU Dioxin Exposure and Health Data - Summary Report*. Retrieved from <http://ec.europa.eu/environment/archives/dioxin/pdf/dioxin.pdf>

EFSA. (2018). *Dioxins and related PCBs: tolerable intake level updated*. Retrieved from <http://www.efsa.europa.eu/en/press/news/dioxins-and-related-pcbs-tolerable-intake-level-updated>

Environment Agency. (2009). *Soil Guideline Values for dioxins, furans and dioxin-like PCBs in soil*. Retrieved from https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/313872/scho0909bqyq-e-e.pdf

Environment Agency. (2016). *Air Emissions Risk Assessment*. Retrieved from <https://www.gov.uk/guidance/air-emissions-risk-assessment-for-your-environmental-permit>

Environmental Protection Scotland and the Royal Town Planning Institute Scotland. (2017). *Delivering Cleaner Air for Scotland*.

European Commission. (2018). *Best Available Techniques (BAT) Reference Document for Waste Incineration*.

HMIP. (1996). *Risk Assessment of Dioxin Releases from Municipal Waste Incineration Processes*.

Hunter and Fewtrell. (2010). *Acceptable Risk*. Retrieved from http://www.who.int/water_sanitation_health/dwq/iwachap10.pdf

Pasternach. (1989). *The Risk Assessment of Environmental and Human Health Hazards*.

Scotland and Northern Ireland Forum for Environmental Research. (2007). *Environmental Legislation and Human Health - Guidance for Assessing Risk*.

WHO. (1998). *Assessment of the Health Risk of Dioxins: Re-evaluation of the Tolerable Daily Intake (TDI)*. Retrieved from <http://www.who.int/ipcs/publications/en/exe-sum-final.pdf>

8 Glossary

2,3,7,8-TCDD	A highly toxic congener
ADMS	Atmospheric Dispersion Modelling System
Congener	An individual dioxin or furan compound
COT	Committee on the Toxicity of Chemicals in Food, Consumer Products and the Environment
Dioxins	polychlorinated dibenzo-para-dioxins
ES	Environmental Statement
Furans	polychlorinated dibenzofurans
Hazard Quotient	The ratio of the predicted exposure level and a 'reference dose' which represents the level at which no adverse effects are expected
HHRAP	Human Health Risk Assessment Protocol
HMIP	Her Majesty's Inspectorate of Pollution
IED	Industrial Emissions Directive
IRAP	Industrial Risk Assessment Program-Human Health
PCB	Polychlorinated Biphenyl
SGV	Soil Guideline Value
TDI	Tolerable Daily Intake
TEQ	International Toxic Equivalence units
TWI	Tolerable Weekly Intake
UK	United Kingdom
USEPA	United States Environmental Protection Agency
WHO	World Health Organisation

Units of Mass

kg	Kilogramme = 1,000 grammes
g	Gramme
µg	Microgramme = 0.000001 grammes
ng	Nanogramme = 0.000000001 grammes
pg	Picogramme = 0.000000000001 grammes

Other units

m³	Cubic metre
Nm³	Normalised cubic metre
s	Second
µm	Micrometer= 0.000001 metres

9 Appendices

A1	Professional Experience.....	27
----	------------------------------	----

A1 Professional Experience

Dr Ben Marner, BSc (Hons) PhD CSci MEnvSc MIAQM

Dr Marner is a Technical Director with AQC and has more than 20 years' experience in the field of air quality. He has been responsible for air quality and greenhouse gas assessments of road schemes, rail schemes, airports, power stations, waste incinerators, commercial developments and residential developments in the UK and abroad. He has been an expert witness at several public inquiries, where he has presented evidence on health-related air quality impacts, the impacts of air quality on sensitive ecosystems, and greenhouse gas impacts. He has extensive experience of using detailed dispersion models, as well as contributing to the development of modelling best practices. Dr Marner has arranged and overseen air quality monitoring surveys, as well as contributing to Defra guidance on harmonising monitoring methods. He has been responsible for air quality review and assessments on behalf of numerous local authorities. He has also developed methods to predict nitrogen deposition fluxes on behalf of the Environment Agency, provided support and advice to the UK Government's air quality review and assessment helpdesk, Transport Scotland, Transport for London, and numerous local authorities. He is a Member of the Institute of Air Quality Management and a Chartered Scientist. Dr Marner is a member of Defra's Air Quality Expert Group.

Dr Imogen Heard, BSc (Hons) MSc PhD MInstPhys

Dr Heard is a Senior Consultant with AQC, having joined the company in 2013. Prior to joining she worked as a scientist in the Atmospheric Dispersion and Air Quality area at the UK Met Office for four years, modelling the dispersion of a range of pollutants over varying spatial and temporal scales. She now works in the field of air quality assessment and has been involved in numerous development projects including road schemes, energy from waste facilities, urban extensions and energy centres. These have included the use of ADMS-5 and ADMS-Roads dispersion models to study the impacts of a variety of pollutants, including nitrogen dioxide, PM₁₀ and PM_{2.5}, and the preparation of air quality assessment reports and air quality chapters for Environmental Statements. She also has experience in undertaking construction dust risk assessments and Air Quality Neutral assessments, as well as in preparing local authority reports.